


Identification and mutation analysis of the complete gene for Chediak-Higashi syndrome

Deborah L. Nagle¹, Mohammad A. Karim², Elizabeth A. Woolf¹, Lisa Holmgren¹, Peer Bork³, Donald J. Misumi¹, Sonja H. McGrail¹, Barry J. Dussault, Jr.¹, Charles M. Perou⁴, Raymond E. Boissy⁵, Geoffrey M. Duyk¹, Richard A. Spritz² & Karen J. Moore¹


Chediak-Higashi syndrome (CHS) is a rare, autosomal recessive disorder characterized by hypopigmentation, severe immunologic deficiency with neutropenia and lack of natural killer (NK) cells, a bleeding tendency and neurologic abnormalities¹⁻⁴. Most patients die in childhood. The CHS hallmark is the occurrence of giant inclusion bodies and organelles in a variety of cell types, and protein sorting defects into these organelles⁵⁻⁸. Similar abnormalities occur in the *beige* mouse^{6,7,9-13}, the proposed model for human CHS. Two groups have recently reported the identification of the *beige* gene^{14,15}, however the two cDNAs were not at all similar. Here we describe the sequence of a human cDNA homologous to mouse *beige*, identify pathologic mutations and clarify the discrepancies of the previous reports. Analysis of the CHS polypeptide demonstrates that its modular architecture is similar to the yeast vacuolar sorting protein, VPS15.

Multiple rounds of screening human cDNA libraries with mouse *beige* probes and subsequently with probes derived from those newly isolated cDNA clones were performed to obtain 27 human cDNA clones. Sequence assembly of these yielded a sequence of 13,449 bp for the human *beige* cDNA homologue, *CHS*, that mapped to chromosome 1q43 by PCR on the G3 Radiation Hybrid Panel¹⁶ (data not shown); a refinement of the previously published map position^{17,18}. A potential translational initiation codon occurs at nt 190, followed by an open reading frame (ORF) of 11,403 bp. A stop codon occurs at nt 11,592 followed by multiple stop codons and a poly(A) tail in a 3'-untranslated region of 1,933 bp.

Fig. 1 Identification of mutations in three patients with CHS. a, Patient 1, a white male with typical childhood CHS, including oculocutaneous albinism (OCA), neutropenia, impaired platelet function, lack of NK cell activity, and characteristic melanosomal abnormalities⁹. Hair and skin are hypopigmented. The patient's parents were third cousins. b, A PCR product spanning codons 461-540 exhibited slightly reduced electrophoretic migration in both the SSCP and duplex patterns, suggestive of a small deletion (data not shown). DNA sequence analysis of the PCR product demonstrated that this patient, shown in a, was homozygous for a single-base deletion within codon 489, resulting in a frameshift distal to this site and translational termination at codon 566. c, Sequence analysis of patient 2 revealed a C-T transition resulting in premature termination and presumptive truncation of the polypeptide at codon 1103. Patient 2, a 27-year old white male with late-onset CHS (lymphoblast culture GM03365; Coriell Institute for Medical Research, Camden, NJ), exhibited albinism, recurrent skin infections, neuropathy and mild mental retardation. No additional clinical information or family history is available.



489
 Lys Ser Glu Gln Leu His
 Normal AAA TCA GAG CAA CTT CAT
 GA-C AAC TTC ATC CCT TGA
 Patient 1 Asp Asn Phe Ile Pro TER



1103
 Ser Ile Arg Leu Leu ...
 Normal ... AGT ATA CGA CTT TTG ...
 Patient 2 ...TGA
 TER

¹ Millennium Pharmaceuticals, Inc., 640 Memorial Dr., Cambridge, Massachusetts 02139, USA
² Departments of Medical Genetics and Pediatrics, University of Wisconsin, Madison, Wisconsin 53706, USA
³ European Molecular Biology Laboratories, Heidelberg, and Max-Delbrück-Centre for Molecular Medicine, Berlin-Buch, Germany
⁴ Division of Cell Biology and Immunology, Department of Pathology, University of Utah School of Medicine, Salt Lake City, Utah 84132, USA
⁵ Department of Dermatology, University of Cincinnati College of Medicine, Cincinnati, Ohio 45267, USA

Correspondence should be addressed to K.M.




Fig. 2 a, Modular architecture of CHS and VPS15. Motifs are described in the text. b, Schematic representation of the position of known mutations (refs. 14, 15 and presented here) along the length of the CHS protein in both mouse (*) and human (◊)

Comparison of *CHS* with the 3' partial 7-kb mouse *beige* cDNA sequence that we reported previously¹⁴ demonstrates 77.2% nucleotide identity and 87.9% amino acid identity between the partial mouse and full-length *CHS* sequences. BLAST searches of the GenBank nucleic acid sequence databases identified multiple expressed sequence tags (ESTs), including those previously identified with the murine *beige* sequence¹⁴. In addition thirteen human ESTs (N25938, H99579, Z21296, Z21358, N39704, W26957, H50968, M78482, H51623, N74354, W03146, N92032, N74383) exhibited almost complete sequence identity to the human *beige* cDNA homologue.

The isolation of the human *CHS* gene allowed us to identify mutations in three patients with CHS (Fig. 1). Patient 1, shown in Fig. 1a, is an inbred boy with typical childhood CHS⁸ and homozygous for a single-base deletion within codon 489, a frameshift mutation that results in premature translational termination at codon 566 (Fig. 1b). Patient 2, an adult male with late-onset CHS, is homozygous for a C to T transition in codon 1103, CGA-to-TGA, resulting in a nonsense mutation (Fig. 1c). Patient 3, a one year-old girl with typical childhood CHS (fibroblast culture GM02075A; Coriell Institute for Medical Research) exhibits 'partial' OCA, photophobia and cytoplasmic inclusions in her white blood cells. She is heterozygous for a previously described frameshift mutation¹⁵; a single-base duplication in codon 40, GCA to GGCA (data not shown). We have not yet identified the second mutation in this patient. The overall similarity of the mouse *beige* and *CHS* gene sequences and the identification of pathologic mutations in patients with CHS definitively prove that *CHS* is homologous to mouse *beige*.

We have recently reported the cloning of the mouse

beige gene¹⁴ as have Barbosa *et al.*¹⁵. However the sequences reported do not align or even partially overlap. Both the partial sequences — that obtained by ourselves and the 4.5-kb sequence obtained by Barbosa *et al.* — are contained within the larger 11,403-bp ORF described here: ours¹⁴ aligns with the 3' end of the ORF of *CHS* while that of Barbosa *et al.* aligns with the 5' end. Both groups have also observed a large 12-kb message in many tissues. The most abundant message observed by Barbosa *et al.* however, was 7 kb. Based upon its abundance in leukocyte cell lines, they concluded that it corresponds with the message of primary functional significance. Nevertheless, as shown in Fig. 2b, pathologic mutations have been identified along the length of the longer, 12-kb mRNA, suggesting that this longer species is critical to function. Surprisingly, the final 36 amino acids of the murine *beige* gene reported by Barbosa *et al.*¹⁵ are not present in the full-length human CHS polypeptide. However, it is possible to PCR-amplify this segment from both mouse genomic DNA and from a bacterial artificial chromosome contig that completely spans the mouse *beige* region (M.D.J. *et al.*, manuscript submitted), suggesting that this sequence represents an alternatively spliced exon.

Additional alternative splice forms of *CHS* have also been isolated; two of the human cDNAs isolated in this study lacked nts 7550–7927. Furthermore, alignment of the murine BG polypeptide sequence reported by Barbosa *et al.*¹⁵ with that of the full-length CHS protein revealed the absence of amino acids 1039–1044 in the former. At present it is not clear whether these differences result from the cloning of splice variants, or from true structural differences between the genes in human and mouse.

merging YSM3_CAEEL and YSM2_CAEEL into one contiguous sequence that appears to contain two genes both related to BG/CHS (Fig. 4). Both significantly match adjacent regions in the portion of CHS that is conserved, and represent two separate loci that are physically linked within the *C. elegans* genome. Of all the ORFs described above, F10F2.1 in *C. elegans* most closely resembles those of *bg* and *CHS*. As much of this sequence homology is restricted to amino acids 3116–3461, we conclude that this is a functionally and structurally defined domain. We have designated this conserved domain BEACH (BEige And CHS) because there is high conservation in otherwise distinct proteins over a wide species range, the length of the region is much larger than protein–protein interaction domains, and because of similarities in predicted alpha-beta folding type and the clustering and properties of the conserved amino acids¹⁴.

The human CHS polypeptide also contains a number of sequence motifs that may provide further clues to its biological function (Fig. 2a). The CHS protein consists of a series of hydrophobic helices, with interspersed hydrophobic regions that do not appear to represent transmembrane domains but are consistent with transmembrane association. These helical regions most closely resemble ARM²⁰ and HEAT²¹ repeat motifs, which tend to form long rods^{20,21}. HEAT repeat motifs occur as a minimum of three repeats in tandem in extremely large proteins that contain extensive helical regions, not unlike CHS. Many of the known HEAT repeat motif-containing proteins are associated with vesicle transport^{22–25}. The C-terminal region of the CHS polypeptide contains seven consecutive WD40 motifs. Four of these precisely correspond with the WD40 consensus, and three have minor deviations²⁶. Such consecutive WD40 motifs form beta sheets arranged in a 7-bladed beta 'propeller fold'^{27,28} that is thought to mediate protein–protein interactions.

Barbosa *et al.*¹⁵ reported that the mouse BG protein exhibits homology to stathmin, a phosphoprotein that regulates the polymerization of microtubules²⁹. We also observe a 26% identity between BG/CHS and stathmin across 72 amino acids (464–536). However, the function of stathmin is dependent on this region forming a coiled-coil, but our analysis of BG/CHS across this region indicates that it has no coiled-coil potential as calculated by using a coiled-coil prediction program³⁰. Furthermore, this small region merely represents 2% of the total size of the protein and comparison of the full-length human CHS polypeptide further weakens the significance of this alignment (1.3×10^{-4} probability by chance). Thus, the significance of the 'homology' between BG/CHS and stathmin remains questionable.

As noted above, the CHS polypeptide shares the closest homology to a number of anonymous ORFs. In addition, the modular architecture of sequence motifs in CHS suggests some potential functions. The presence of multiple hydrophobic regions resembling related ARM and HEAT repeats¹⁹, as well as 4–7 consecutive WD40 sequence motifs, suggests that CHS may be a cytoplasmic protein involved in transport and/or associated with vesicles. The only known protein that contains HEAT repeats (or helical regions that resemble HEAT and ARM repeats), C-terminal consecutive WD40 motifs and a globular alpha/beta domain is the yeast serine/threonine

protein kinase, VPS15. *Vps15* is a member of a large class of yeast 'Vps' mutants that are associated with defective vacuolar protein sorting, and fall into more than 40 complementation groups (reviewed in ref. 31). *Vps* mutants exhibit defective sorting in that they tend to secrete soluble vacuolar hydrolase precursors instead of sorting them into the vacuolar compartment³². It is thought that VPS15 is required for activation of a second VPS (VPS34) and that they function together as components of a membrane-associated signal transduction complex that regulates intracellular protein trafficking³³.

Given the similarity in modular architecture of VPS15 and the BG/CHS proteins (Fig. 2a), we suggest that the BG/CHS proteins may have a similar function, consistent with the observation of defective vesicular transport to and from the lysosome and late endosome^{6,8,10,12} and aberrant compartmentalization of lysosomal and granular enzymes^{5,7,8,10,11} both in humans with CHS and in mice carrying *beige* mutations. The role of BG/CHS as a component of a membrane-associated signal transduction complex that regulates intracellular protein trafficking is also consistent with our observations that multiple BG/CHS paralogues exist in *C. elegans*. This suggests that BG/CHS may define a novel gene family; an hypothesis compatible with both the existence of multiple mouse loci, mutants of which have phenotypes that are partially similar to *beige* mutants³⁴, and with the potential of CHS being a heterogeneous disease¹⁷.

Methods

CHS cDNA isolation. To isolate the complete human *beige* cDNA homologue, we first screened human fetal liver and retina cDNA libraries (Clontech) with probes from multiple regions of the murine *beige* cDNA¹⁴. Subsequently, we carried out multiple rounds of screening using probes derived from the newly isolated human *beige* homologous cDNAs. In total, 27 cDNAs (12 from a fetal liver library and 15 from a human retina library) were identified and sequenced completely on both strands.

Genetic mapping. Sequence-tagged sites (STSs) were designed to multiple segments along the length of the cDNA; all STSs were mapped by PCR on the G3 Radiation Hybrid Panel¹⁶.

Mutation detection. High molecular weight genomic DNA was prepared from cultured cells from patients with CHS by standard methods. DNA segments were amplified from genomic DNAs of the patients and at least two normal individuals by PCR using primers derived from the cDNA sequence, and analysed by simultaneous single-strand conformation polymorphism (SSCP)/heteroduplex (HDX) analyses. Segments exhibiting aberrant SSCP/HDX patterns were either reamplified from each patient in duplicate, and cloned into pCR2.1 (Invitrogen, San Diego, CA) from which at least six independent clones were sequenced, or sequenced directly from multiple independent PCR amplifications.

Structural analysis. Initial database searches were performed using the BLAST series of programs³⁵. Putative nonglobular domains that function as linkers were identified using the SEG program (ref. 36 and refs therein). Secondary structure prediction and screening for transmembrane helices were carried out using the PHD Web server implement³⁷. The coiled-coil potential was measured using the COILS program³⁰. To increase the sensitivity of the database similarity search programs, CHS was partitioned into segments according to similar secondary struc-

ture content and to putative globular domains. As homologous sequences were identified, iterate motif and profile searches were performed³⁸.

GenBank accession number. The accession number for the CHS cDNA sequence is U67615.

Acknowledgements

We thank N. Lakey and members of the Sequencing group at MPI, and J. Kaplan for helpful discussions.

Received 4 September; accepted 19 September 1996.

- Beguez-Cesar, A.B. Neutropenia cronica maligna familiar con granulaciones atipicas de los leucocitos. *Bol. Soc. Cubana Pediatr.* **15**, 900-922 (1943).
- Steinbrinck, W. Über eine neue Granulationsanomalie der Leukocyten. *Dtsch. Arch. Klin. Med.* **193**, 577-581 (1948).
- Chediak, M. Nouvelle anomalie leucocytaire de caractere constitutionnel et familial. *Rev. Hematol.* **7**, 362-367 (1952).
- Higashi, O. Congenital gigantism of peroxidase granules. *Tohoku J. Exp. Med.* **59**, 315-332 (1954).
- Jones, K.L., Stewart, R.M., Fowler, M., Fukuda, M. & Holcombe, R.F. Chediak-Higashi lymphoblastoid cell lines: granule characteristics and expression of lysosome-associated membrane proteins. *Clin. Immunol. Immunopath.* **65**, 219-226 (1992).
- Burkhardt, J.K., Wiebel, F.A., Hester, S. & Argon, Y. The giant organelles in Beige and Chediak-Higashi fibroblasts are derived from late endosomes and mature lysosomes. *J. Exp. Med.* **178**, 1845-1856 (1993).
- Holcombe, R.F., Jones, K.L. & Stewart, R.M. Lysosomal enzyme activities in Chediak-Higashi syndrome: evaluation of lymphoblastoid cell lines and review of the literature. *Immunodeficiency* **5**, 131-140 (1994).
- Zhao, H. *et al.* On the analysis of the pathophysiology of Chediak-Higashi syndrome. *Lab. Invest.* **71**, 25-34 (1994).
- Lutzner, M.A., Lowrie, C.T. & Jordan, H.W. Giant granules in leukocytes of the beige mouse. *Heredity* **58**, 299-300 (1966).
- Brandt, E.J., Elliott, R.W. & Swank, R.T. Defective lysosomal enzyme secretion in kidneys of Chediak-Higashi (beige) mice. *J. Cell Biol.* **67**, 774-788 (1975).
- Swank, R.T. & Brandt, E.J. Turnover of kidney β -glucuronidase in normal and Chediak-Higashi (beige) mice. *Am. J. Pathol.* **92**, 755-771 (1978).
- Willingham, M.C., Spicer, S.S. & Vincent, R.A., Jr. The origin and fate of large dense bodies in beige mouse fibroblasts. *Exp. Cell Res.* **136**, 157-168 (1981).
- Penner, J.D. & Prieur, D.J. A comparative study of the lesions in cultured fibroblasts of humans and four species of animals with Chediak-Higashi syndrome. *Am. J. Med. Genet.* **28**, 445-454 (1987).
- Perou, C.M. *et al.* Identification of the murine beige gene by YAC complementation and positional cloning. *Nature Genet.* **13**, 303-308 (1996).
- Barbosa, M.D.F.S. *et al.* Identification of the homologous beige and Chediak-Higashi syndrome genes. *Nature* **382**, 262-265 (1996).
- Stein, L., Kruglyak, L., Slonim, D. & Lander, E. Unpublished software, Whitehead Institute/MIT Center for Genome Research (1995).
- Fukai, K. *et al.* Homozygosity mapping of the gene for Chediak-Higashi syndrome to chromosome 1q42-q44 in a segment of conserved synteny that includes the mouse beige locus (bg). *Am. J. Hum. Genet.* **59**, 620-624 (1996).
- Barret, F.J. *et al.* Genetic and physical mapping of the Chediak-Higashi syndrome on chromosome 1q42-43. *Am. J. Hum. Genet.* **59**, 625-632 (1996).
- Bork, P. & Koonin, E.V. Protein sequence motifs. *Curr. Opin. Struct. Biol.* **6**, 366-376 (1996).
- Peifer, M., Berg, S. & Reynolds, A.B. A repeating amino acid motif shared by proteins with diverse cellular roles. *Cell* **76**, 789-791 (1994).
- Andrade, M.A. & Bork, P. HEAT repeats in the Huntington's disease protein. *Nature Genet.* **11**, 115-116 (1995).
- DiFiglia, M. *et al.* Huntingtin is a cytoplasmic protein associated with vesicles in human and rat brain neurons. *Neuron* **14**, 1075-1081 (1995).
- Sabatini, D.M., Ercjument-Bromage, H., Lui, M., Tempst, P. & Snyder, S.H. RAFT1: a mammalian protein that binds to FKBP12 in a rapamycin-dependent fashion and is homologous to yeast TORs. *Cell* **78**, 35-43 (1994).
- Zheng, X. F., Florentino, D., Chen, J., Crabtree, G. R. & Schreiber, S. L. TOR kinase domains are required for two distinct functions, only one of which is inhibited by rapamycin. *Cell* **82**, 121-130 (1995).
- Hemmings, B. A. *et al.* α - and β -forms of the 65-kDa subunit of protein phosphatase 2A have a similar 39 amino acid repeating structure. *Biochemistry* **29**, 3166-3173 (1990).
- Neer, E.J., Schmidt, C.J., Nambudripad, R. & Smith, T.F. The ancient regulatory-protein family of WD-repeat proteins. *Nature* **371**, 297-300 (1994).
- Sondek, J., Bohm, A., Lambright, D. G., Hamm, H. E. & Sigler, P. B. Crystal structure of a G α protein beta gamma dimer at 2.1 Å resolution. *Nature* **379**, 369-374 (1996).
- Wall, M. A. *et al.* The structure of the G protein heterotrimer G β γ 2. **83**, 1047-1058 (1995).
- Belmont, L. D. & Mitchison, T. J. Identification of a protein that interacts with tubulin dimers and increases the catastrophe rate of microtubules. *Cell* **84**, 623-631 (1996).
- Lupas, A.N. *et al.* Predicting coiled coils from protein sequences. *Science* **252**, 1162-1164 (1991).
- Klionsky, D. J. & Emr, S. D. A new class of lysosomal/vacuolar protein sorting signals. *J. Biol. Chem.* **265**, 5349-5352 (1990).
- Herman, P. K., Stack, J. H. & Emr, S. D. A genetic and structural analysis of the yeast Vps15 protein kinase: evidence for a direct role of VPS15p in vacuolar protein delivery. *EMBO J.* **10**, 4049-60 (1991).
- Stack, J. H., Herman, P. K., Schu, P. V. & Emr, S. D. A membrane-associated complex containing the Vps15 protein kinase and the VPS34 PI 3-kinase is essential for protein sorting to the yeast lysosome-like vacuole. *EMBO J.* **12**, 2195-204 (1993).
- Novak, E.K., Hui, S.W. & Swank, R.T. Platelet storage pool deficiency in mouse pigment mutations associated with seven distinct genetic loci. *Blood* **63**, 536-544 (1984).
- Altschul, S. F., Boguski, M. S., Gish, W. & Wootton, J. C. Issues in searching molecular sequence databases. *Nature Genet.* **6**, 119-129 (1994).
- Wootton, J. C. & Federhen, S. Analysis of compositionally biased regions in sequence databases. *Meth. Enz.* **266**, 554-571 (1996).
- Rost, B., Sander, C. & Schneider, R. PHD—an automatic mail server for protein secondary structure prediction. *Comput. Appl. Biosci.* **10**, 53-60 (1994).
- Bork, P. & Gibson, T. Applying motif and profile searches. *Meth. Enz.* **266**, 162-184 (1996).